

**GUIDELINE FOR THE MANAGEMENT OF TENSION PNEUMOTHORAX IN
NEONATES**

This is the most current document and should be used until a revised version is available

Key Document code:	WAHT-KD-015	
Key Documents Owner:	Dr Vivianna Weckemann	Consultant Paediatrician
Approved by:	Paediatric Quality Improvement Meeting	
Date of Approval:	2 nd August 2019	
Date of Review:	2 nd August 2022	
This is the most current version and should be used until a revised document is in place		

Key Amendments

Date	Amendment	Approved By

INTRODUCTION


Tension pneumothorax is a rare, potentially fatal condition primarily occurring in very sick, ventilated infants. Following the widespread use of surfactant therapy and changes in ventilation strategies to avoid volutrauma, occurrence of Pneumothorax has dropped but prevention is there fore essential. The continued use of antenatal steroids, early natural surfactants and ventilation regimes that avoid excessive minute volumes and tidal volumes are other preventative measures.

GUIDELINE

Signs of Tension Pneumothorax

- Sudden deterioration in the infant with worsening respiratory distress and rising oxygen requirements/falling oxygen saturation.
- Hyper-resonance with decreased air entry on affected side.
- Shift in position of cardiac impulse/heart sounds.
- Abdominal distension.
- Positive transillumination on affected side.

<p>Please note that the key documents are not designed to be printed, but to be used on-line. This is to ensure that the correct and most up-to-date version is being used. If, in exceptional circumstances, you need to print a copy, please note that the information will only be valid for 24 hours and should be read in conjunction with the key document supporting information and/or Key Document intranet page, which will provide approval and review information.</p>
--



Treatment

If the baby is stable, and not deteriorating, arrange urgent chest X-ray to confirm diagnosis.

If situation is critical and transillumination positive, insert formal chest drain* on affected side.

If situation critical and transillumination equivocal with a high index of suspicion perform needle aspiration with 21g butterfly needle, 10ml. syringe and 3 ways tap at 2nd intercostals space in the mid clavicle line.

We stock two types of chest drain at present until all staff are comfortable with the seldinger technique drains as these are the preferred type of drain on NNU.

****Technique for Formal Chest Drain Insertion- using Vygon “conventional” chest drainage***

Equipment needed

- sterile dressing pack
- scalpel and straight surgical blade
- artery forceps
- suture (3.0 silk)
- intercostal drain (10 FG)
- connecting pigot and Heimlich valve – this comes with the chest drain in the same box



Procedure

1. Ensure adequate analgesia (Morphine IV ideally).
2. Position baby with affected side elevated and arm on affected side elevated.
3. Clean skin over mid axillary line and infiltrate 1% lignocaine at insertion site over 5th/6th intercostal space well away from breast bud.
4. Make small incision down to intercostal muscles using scalpel blade.
5. Use artery forceps for blunt dissection down to pleura.
6. Remove trochar from the chest drain.
7. Grasp tip of drain with artery forceps and insert through pleural sac, listening for gas leak - insert to depth of 2-3 cm.
8. Connect drain to Heimlich valve using appropriate connector.
9. Suture the drain in place and anchor with adhesive tape.
10. Perform chest X-ray to confirm position of tube and resolution of pneumothorax.



*Technique for Formal Chest Drain Insertion- using Cook “Seldinger technique” chest drainage

NB there is an opened set kept in NNU intensive care nursery for staff to use to familiarise themselves with the kit

Equipment needed

- sterile dressing pack
- scalpel and straight surgical blade
- suture (3.0 silk)
- intercostal drain (10 FG)
- connecting spigot and Heimlich valve – this comes in a separate packet to the chest drain




Chest Drain



flutter valve


WAHT-KD-015
Neonatal Key DocumentsProcedure

1. Ensure adequate analgesia (Morphine IV ideally).
2. Position baby with affected side elevated and arm on affected side elevated.
3. Clean skin over mid axillary line and infiltrate 1% lignocaine at insertion site over 5th/6th intercostal space well away from breast bud.
4. Make small incision of a few mm in the skin using scalpel blade.
5. Pass the needle from the chest drain kit through the incision into the pleural cavity



6. Pass the guide wire down the needle and then withdraw the needle leaving the guide wire in situ






7. Pass the dilator over the guide wire and pass it down along the wire and through the chest wall so that the tip is within the pleural space. This dilates the track for easy passage of the drain. Now remove the dilator leaving the wire in situ.



8. Pass the chest drain over the wire and through the chest wall ensuring that all drainage holes are within the pleural space.
9. Remove the guide wire leaving the chest drain in situ
10. Connect drain to the Heimlich flutter valve use the blue set that is made by Cook medical
11. Secure drain using suture (not purse string) and clear adhesive dressing



WAHT-KD-015
Neonatal Key Documents

MONITORING TOOL

How will monitoring be carried out? Individual case review

Who will monitor compliance with the guideline? Consultant staff

STANDARDS:

Item	%	Exceptions
Compliance with insertion technique of chest drains on all babies meeting criteria	100%	none

REFERENCES

Textbook of Neonatology Ed. Rennie + Robertson 3rd. Ed. pp. 517-521; 1381-1382.

Please note that the key documents are not designed to be printed, but to be used on-line. This is to ensure that the correct and most up-to-date version is being used. If, in exceptional circumstances, you need to print a copy, please note that the information will only be valid for 24 hours and should be read in conjunction with the key document supporting information and/or Key Document intranet page, which will provide approval and review information.

CONTRIBUTION LIST

Key individuals involved in developing the document

Name	Designation
Dr Andrew Short	Consultant Paediatrician WRH

Circulated to the following individuals for comments

Name	Designation
Dr A Gallagher	Consultant Paediatrician, WRH
Dr D.Castling	Consultant Paediatrician ,WRH
Dr V Weckemann	Consultant Paediatrician ,WRH
Dr M.Hanlon	Consultant Paediatrician ,WRH
Dr J.Scanlon	Consultant Paediatrician ,WRH
Dr L Harry	Consultant Paediatrician, WRH
Dr C Onyon	Consultant Paediatrician ,ALEX
Dr K.Nathavitharana	Consultant Paediatrician ,ALEX
Dr N.Ahmad	Consultant Paediatrician ,ALEX
Dr M. Ahmed	Consultant Paediatrician, ALEX
Dr T Dawson	Consultant Paediatrician, ALEX
Vicky Bullock	Matron NICU
Karen Kokoska	Maternity Risk Manager
Patti Paine	Head of Midwifery
Margaret Stewart	Matron OP/Community
Alison Talbot	Matron, Maternity IP, Alex
Rachel Carter	Matron, Maternity IP, WRH
Matt Kaye	Clinical Pharmacist

Circulated to the following CD's/Heads of dept for comments from their directorates / departments

Name	Directorate / Department

Circulated to the chair of the following committee's / groups for comments

Name	Committee / group
Alison Smith	Medicines Safety Committee