

## Ketamine infusion for pain relief in Adult patients

<b>Key Document Code:</b>	WAHT-KD-004
<b>Key Document Owner:</b>	Dr Hutchinson
<b>Approval Date:</b>	12 <sup>TH</sup> November 2019
<b>Review Date:</b> <b>This is the most current document and should be used until a revised version is in place</b>	12 <sup>th</sup> November 2022

### Key Amendments

Date	Amendment	Approved by

### Introduction

The majority of postoperative pain is controlled using simple analgesics, opioids or central neuraxial blockade. Opioids are known to produce tolerance and hyperalgesia. Ketamine is an anaesthetic agent with analgesic properties in sub anaesthetic doses. Ketamine exerts its main analgesic effect by antagonism of NMDA receptors and modulates central sensory processing of pain. Ketamine has been shown to have a potent antihyperalgesic effect and can counteract opioid-induced hyperalgesia and prevent the development of opioid tolerance.<sup>(1)</sup> Ketamine infusions can be used in certain circumstances such as surgery associated with high analgesic requirements, refractory neuropathic pain, existing opioid tolerance, allergy to opioids or intolerance of side effects.

### Details of Guideline

Ketamine infusions should only be used in the Critical care units or Surgical and Vascular High Dependency Units. Ketamine infusions should be prescribed on an "Injectable Medicines" prescription chart.

A small dose of intravenous ketamine usually 2.5-5mg and no more than 10mg may be administered by an Anaesthetist or member of the Acute Pain team to assess efficacy before commencing an infusion. It is important to inform the patient that they may feel a little strange before administration of the test dose. Most patients notice a spaced out sensation which is tolerated much better if they are pre-warned. If judged to be effective and well tolerated, an infusion is commenced.

200mg ketamine should be made up to 40ml using 0.9% Saline, giving a concentration of 5mg/ml.  
Infusion rate is between 0-3ml/hr

### Contraindications

#### Absolute Contraindications:

Allergy to ketamine  
Acute Porphyria

#### Relative Contraindications

Ischaemic heart disease  
Uncontrolled hypertension  
Raised intracranial pressure  
Raised intraocular pressure  
Psychosis

#### Side Effects

Hypertension  
Tachycardia  
Diplopia

Please note that the key documents are not designed to be printed, but to be used on-line. This is to ensure that the correct and most up-to-date version is being used. If, in exceptional circumstances, you need to print a copy, please note that the information will only be valid for 24 hours and should be read in conjunction with the key document supporting information and/or Key Document intranet page, which will provide approval and review information.

Nystagmus  
Rash  
Nausea and Vomiting  
Hallucinations  
Hypersalivation

At higher doses sedation may occur. This is more common when ketamine is prescribed with other sedative agents eg. high doses of opioids, anticonvulsants and benzodiazepines.

Patients may experience a sense of dissociation from reality, dysphoria and rarely vivid dreams or hallucinations. If these symptoms occur, reducing the infusion rate may be effective whilst still providing analgesia. If symptoms persist, discontinue the infusion.